

# Foot and Mouth Disease outbreak report

---

2021-2023



agriculture, land reform  
& rural development

---

Department:  
Agriculture, Land Reform and Rural Development  
**REPUBLIC OF SOUTH AFRICA**

Report compiled by:  
Directorate: Animal Health

# 1. Introduction and summary


South Africa currently has three Foot and Mouth Disease (FMD) outbreak events in the previously FMD free zone without vaccination. These three outbreak events currently comprise 200 open outbreaks<sup>1</sup> reported to the World Organisation for Animal Health (WOAH founded as OIE). South Africa has so far resolved 12 infected premises in the three FMD outbreak events and closed them with the WOAH.

The first outbreak event started in May 2021 and mainly affects KwaZulu-Natal (KZN) Province, with limited spread to Free State Province. The second outbreak event started in March 2022 in the previous FMD free zone in Limpopo Province, with limited spread to Gauteng Province. The third outbreak event also started in March 2022 in North West Province (linked to the second outbreak event in Limpopo Province) with spread to Mpumalanga, Gauteng and Free State Provinces.

There are two viruses responsible for the current three outbreak events:

- SAT 2 serotype in outbreak event 1 (KwaZulu-Natal – Free State)
- SAT 3 serotype in outbreak event 2 (Limpopo – Gauteng) and outbreak event 3 (North West – Gauteng – Mpumalanga – Free State)

**Map 1: Reported outbreaks in the previous FMD free zone**



<sup>1</sup>'Outbreak' means the occurrence of one or more cases in an epidemiological unit. An 'event' means a single outbreak or a group of epidemiologically related outbreaks of a given disease.

**Table 1: Summary of outbreaks per outbreak event and per Province:**

Outbreak Event	Province	Number of open outbreaks	Number of resolved outbreaks	Total number of outbreaks	Start date of last reported outbreak
Outbreak event 1: KwaZulu-Natal – Free State	KZN	136	2	138	18 Feb 2023
	FS	3	1	4	26 Sep 2022
Outbreak event 2: Limpopo - Gauteng	LP	7	1	8	25 Apr 2022
	GP	0	2	2	30 Mar 2022
Outbreak event 3: North West – Gauteng – Mpumalanga – Free State	NW	16	2	18	22 Nov 2022
	GP	4	1	5	30 Aug 2022
	MP	1	0	1	2 Aug 2022
	FS	33	3	36	22 Sep 2022
<b>Total</b>		<b>200</b>	<b>12</b>	<b>212</b>	

## **2. Details of open outbreak events (as reported to the WOA)**

### **2.1 Outbreak event 1: KwaZulu-Natal – Free State Provinces**


This outbreak event involves 139 locations in total that are currently still open with the WOA. This comprises 136 locations in KZN which are mainly communal dip tanks and a few small-scale farms. In the past month, since the previous update report of 28 March 2023, there were 20 positive cases reported in KZN. Most of these cases occurred in 2022, with a few in January and February 2023, but were only reported now, and the clinical picture on the ground does not indicate current spread.

Following spread of this outbreak to the adjacent Free State province in September 2022, there are still 3 open outbreaks in the Free State, including one feedlot and two small scale farms. However, no further spread was detected in the Free State in the past 7 months. The remaining affected farms are still under quarantine.

Three outbreaks in this event have been resolved and closed with the WOA. This includes two feedlots in KZN, with both feedlots depopulated through controlled slaughter. In the Free State, one outbreak occurred in a communal area adjacent to a feedlot. This outbreak was resolved by absorbing the animals into the affected feedlot.

In June 2022, FMD positive buffalo were detected in a wildlife reserve in KZN and towards the end of 2022, buffalo in another game reserve in the area had also become infected. Since it may not be possible to destroy all the infected buffalo populations, the carrier status of buffalo for FMD will mean that the affected area will remain under FMD control.

**Map 2: Outbreak event 1 in KwaZulu-Natal – Free State Provinces**




## **2.2 Outbreak event 2: Limpopo – Gauteng Province**

In the past month, since the previous update report of 28 March 2023, the situation in Limpopo Province remained stable. The start date of the last outbreak in the Province was 25 April 2022, which marks a 12 month period with no new outbreaks reported.

7 outbreaks in communal diptanks remain open with the WOAH. Clinical and serological surveillance is underway to confirm the absence of circulating virus, following which the outbreak event will be closed with the WOAH.

In April 2022, serologically positive FMD vaccinated animals were found at two premises in Gauteng Province and one in Limpopo Province. Investigations identified that these animals had been illegally moved from the Protection Zone with vaccination in Limpopo Province. The animals were confiscated, and safely moved to a designated abattoir in Limpopo Province for controlled slaughter. These three outbreaks in this event were thus closed with WOA in May and July 2022.

**Map 3: Outbreak event 2 in Limpopo - Gauteng Provinces**



### **2.3 Outbreak event 3: North West - Free State – Gauteng - Mpumalanga Provinces**

In the past month, since the previous update report of 28 March 2023, the situation in this outbreak event remained stable. There are currently 54 open outbreaks in this event, involving 4 provinces, while 6 outbreaks have been resolved and closed with the WOA.


In the North West Province, the start date of the last outbreak was more than 5 months ago. Two affected farms – a feedlot and a commercial farm - have been depopulated through controlled slaughter and resolved with the WOA. Sixteen (16) farms, comprising commercial breeding herds and stud farms, remain under quarantine.

In the Free State province, 33 commercial farms remain under quarantine and no new outbreaks have been reported in the past 7 months. Three (3) farms have been resolved through controlled slaughter and closed with the WOAH.

In Gauteng province, 4 properties remain under quarantine, with the start date of the last outbreak more than 8 months ago. One affected farm was depopulated through controlled slaughter and the outbreak closed with the WOAH. Another affected feedlot has been depopulated through controlled slaughter and will be closed with the WOAH as soon as disinfection has concluded.

In Mpumalanga, one feedlot was affected and no new outbreaks were reported in the past 9 months. The feedlot was effectively depopulated through controlled slaughter and the outbreak will be closed with the WOAH once quarantine is lifted.

**Map 4: Outbreak event North West - Free State – Gauteng – Mpumalanga Provinces**



### **3. Control Measures**

#### **3.1 Movement control**

##### **3.1.1 Quarantine of affected premises:**

All affected farms and feedlots in North West, Free State, Gauteng and Mpumalanga Provinces were placed under quarantine with strict movement control. The locations involved are well fenced and movement of animals from these farms can be effectively prevented.

##### **3.1.2 Declaration of Disease Management Areas:**

Where the outbreaks occurred in communal grazing land in KZN Province and Limpopo Province, the Minister of Agriculture, Land Reform and Rural Development declared Disease Management Areas (DMA) in the Government Gazette to cover the wider epidemiologically high-risk areas. A DMA was also declared in the Free State Province to include all areas with positive locations or perceived to be at high risk of infection due to being epidemiologically linked.

With the three DMAs, cattle, sheep and goats were not allowed to be moved into, out of and through the area, as well as from one property to another for any reason, except for direct slaughter at a registered abattoir. Such movements required veterinary health certification and a movement permit. Cattle, sheep and goats were allowed to be slaughtered for own consumption or for cultural or religious purposes, but only in the same epidemiological area where they were kept prior to slaughter. Visible Veterinary Patrols and roving and stationary road blocks were directed to cover high risk areas according to information on possible movement of animals.

The Free State DMA has since been lifted, while the KZN and Limpopo DMAs were made smaller based on risk assessment and surveillance of the areas involved. The movement of cattle, sheep and goats for purposes other than direct slaughter will be considered from FMD negative properties in these areas, if deemed low risk and under specific risk mitigation conditions.

### 3.1.3 National movement ban:

During 2022, the outbreaks within the country were spreading despite the control measures implemented. A national movement ban on cattle was instituted from 16 August 2022 until 8 September 2022. During this period, cattle could not be moved from one property to another for any reason, except for direct slaughter at a registered abattoir. No cattle could be moved between farms, locations or premises for purposes of sale, shows, breeding, backgrounding, ritual purposes, lobola, or any other purpose. No cattle could be moved into feedlots, and movement out of feedlots were only allowed directly to registered abattoirs. No cattle could be moved for the purpose of import or export.

### 3.1.4 Movement control in unaffected locations:

Since the upliftment of the national movement ban, all livestock owners must provide a declaration of health and origin and must obtain the necessary Stock Theft documents for all cattle, sheep and goats to be moved from their property of origin onto other properties. Recipients of such animals must give written agreement to isolate new livestock for 28 days before introducing them into the main herds of the destination farm. The control measures for movement of cloven-hoofed animals and products out of the FMD protection zones have not changed.

## **3.2 Vaccination**

Approximately 634 000 animals have been vaccinated to control the FMD outbreaks. Vaccination was implemented in March 2022 to curtail the spread of the disease. Cattle in communal areas are vaccinated on a risk-based approach, to ensure that the areas at highest risk are vaccinated first. Cattle on affected farms and feedlots were also vaccinated to reduce the viral load and lower the risk of spreading of disease while the controlled slaughter processes are ongoing.

Routine vaccination in the protection zone is also continuing. The vaccine being used is procured from the Botswana Vaccine Institute and encompasses all three SAT types. Only government officials are allowed to vaccinate animals as per the outbreak control plans.



### **3.3 Controlled slaughter**

Affected positive feedlots and certain commercial farms are depopulated through controlled slaughter at designated abattoirs. The risk materials are processed or disposed of and only known safe products are released into the local market. No meat from affected animals may enter the export market. The infected properties remain under quarantine until 28 days after depopulation and disinfection. The outbreaks on the properties are officially closed once the disinfection process has been concluded. Around 200 000 animals have been sent for controlled slaughter to date.

## **4. Confirmation of diagnosis**

Confirmation of disease was done using a combination of the Solid Phase Competition ELISA (SPCE), Non Structural Protein (NSP) ELISA, and Polymerase Chain Reaction (PCR). All diagnostic tests are performed at the ARC Onderstepoort Veterinary Research Transboundary Animal Diseases laboratory (OVR-TAD).

## **5. Epidemiological investigation**

The SAT 3 virus, responsible for outbreak events 2 and 3, originated in Limpopo Province. The source of introduction of this virus into Limpopo could not be determined, as the virus is not closely linked to any viruses recently identified in the infected or protection zones of South Africa.


This virus spread to North West Province, and subsequently to the Free State, Gauteng and Mpumalanga Provinces. The animals that caused the outbreak in North West Province moved from an area in Limpopo Province that was not under restriction at the time of the movement.

In KwaZulu Natal Province, epidemiological investigations to date have not revealed a source for the outbreak. However, the virus responsible for the outbreak is a SAT 2 serotype and is closely related to a SAT 2 virus responsible for an outbreak that occurred in the Protection Zone in northern Limpopo Province in 2019. It is therefore most likely that illegal movement of animals from the Protection Zone in Limpopo to KZN Province, may be the cause of the outbreak in KZN.

## 6. Surveillance

Clinical and serological surveillance around the affected areas continue, with intensified inspections around identified infected dip tanks in communal areas, or epidemiologically linked locations identified through forward and backward tracing. The clinical presentation of the disease in different locations have varied, from the classical clinical picture in some locations, to absence of clinical signs in others. Surveillance is therefore based on both clinical inspections, including mouthing, as well as serology.

All cattle presented at auctions are inspected before being admitted to the auction pens. If any animal presents with clinical signs, the auction is suspended while investigations are conducted, including laboratory confirmation ruling out FMD.

  
\_\_\_\_\_  
**Director Animal Health**  
Date: 2023-04-30